



ELSEVIER

Contents lists available at ScienceDirect

## Developmental and Comparative Immunology

journal homepage: [www.elsevier.com/locate/devcompimm](http://www.elsevier.com/locate/devcompimm)

# Manipulation of the silkworm immune system by a metalloprotease from the pathogenic bacterium *Pseudomonas aeruginosa*

Li Ma<sup>a</sup>, Lizhen Zhou<sup>a</sup>, Jinshui Lin<sup>b</sup>, Jiuyuan Ji<sup>a</sup>, Yang Wang<sup>c</sup>, Haobo Jiang<sup>c</sup>, Xihui Shen<sup>b</sup>, Zhiqiang Lu<sup>a,\*</sup>

<sup>a</sup> Department of Entomology, College of Plant Protection, Northwest A&F University, Yangling, Shaanxi 712100, China

<sup>b</sup> Department of Microbiology, College of Life Science, Northwest A&F University, Yangling, Shaanxi 712100, China

<sup>c</sup> Department of Entomology and Plant Pathology, Oklahoma State University, Stillwater, OK, 74078, USA

## ARTICLE INFO

## Keywords:

*Bombyx mori**Pseudomonas aeruginosa*

Immune

Elastase B

Phenoloxidase

Antimicrobial peptide

## ABSTRACT

Antimicrobial peptide (AMP) production and melanization are two key humoral immune responses in insects. Induced synthesis of AMPs results from Toll and IMD signal transduction whereas melanization depends on prophenoloxidase (PPO) activation system. During invasion, pathogens produce toxins and other virulent factors to counteract host immune responses. Here we show that the pathways leading to PPO activation and AMP synthesis in the silkworm *Bombyx mori* are affected by a metalloprotease, named elastase B, secreted by *Pseudomonas aeruginosa* (PAO1). The metalloprotease gene (*lasB*) was expressed shortly after PAO1 cells had been injected into the larval silkworm hemocoel, leading to an increase of elastase activity. Injection of the purified PAO1 elastase B into silkworm hemolymph compromised PPO activation. In contrast, the protease caused a level increase of gloverin, an AMP in the hemolymph. To verify our results obtained using the purified elastase B, we infected *B. mori* with PAO1  $\Delta lasB$  mutant and found that PO activity in hemolymph of the PAO1  $\Delta lasB$ -infected larvae was significantly higher than that in the wild type-infected. The mutant-inhabited hemolymph had lower levels of gloverin and antimicrobial activity. PAO1  $\Delta lasB$  showed a decreased viability in the silkworm hemolymph whereas the host had a lower mortality. In addition, the effects caused by the  $\Delta lasB$  mutant were restored by a complementary strain. These data collectively indicated that the elastase B produced by PAO1 is an important virulent factor that manipulates the silkworm immune system during infection.

## 1. Introduction

Insects rely on an innate immune system to defend themselves against pathogenic bacteria. This system consists of cellular and humoral responses. The former is mediated by hemocytes including phagocytosis, nodulation and encapsulation (Strand, 2008) and the latter includes melanization and antimicrobial peptide (AMP) synthesis. Melanization is catalyzed by phenoloxidase (PO) produced in a prophenoloxidase (PPO) activation system. AMP synthesis is induced by NF- $\kappa$ B signaling pathways (Vallet-Gely et al., 2008). Similar to other insects, the silkworm *Bombyx mori* employs POs and AMPs to cope with bacterial infection (Tanaka et al., 2008; Huang et al., 2009; Yang et al., 2011).

To avoid the attacks by the host immune system, bacteria may hide

from detection or recognition (Hurst et al., 2003). In other cases, bacteria may suppress the host defense responses in various manners. For instance, an entomopathogenic bacterium *Photobacterium luminescens* produces an antibiotic (E)-1,3-dihydroxy-2-(isopropyl)-5-(2-phenylethenyl) benzene that inhibits phenoloxidase (PO) of the tobacco hornworm *Manduca sexta* (Eleftherianos et al., 2007). Live *Xenorhabdus hominickii* ANU101 infection reduces PO activity and inhibits AMP expression in the beet armyworm *Spodoptera exigua* (Sadekuzzaman et al., 2017). *Serratia marcescens* infection induces apoptosis of the silkworm hemocytes (Ishii et al., 2012). A metalloprotease serralyisin secreted by *S. marcescens* disrupts the adhesive properties of hemocytes and suppresses wound healing, leading to a massive loss of hemolymph in the silkworm (Ishii et al., 2014a, 2014b).

The Gram-negative bacterium *P. aeruginosa* is a widespread

**Abbreviations:** AMP, antimicrobial peptide; BSA, bovine serum albumin; CFU, colony forming units; HPLC, high-performance liquid chromatography; IMD, immune deficiency; LC-MS/MS, liquid chromatography-tandem mass spectrometry; LB, Luria-Bertani; PBS, phosphate-buffered saline; PO, phenoloxidase; PPO, prophenoloxidase; SDS-PAGE, sodium dodecyl sulfate polyacrylamide gel electrophoresis; TCA, trichloroacetic acid

\* Corresponding author.

E-mail address: [zhiqiang.lu@nwsuaf.edu.cn](mailto:zhiqiang.lu@nwsuaf.edu.cn) (Z. Lu).

<https://doi.org/10.1016/j.dci.2018.09.017>

Received 12 August 2018; Received in revised form 22 September 2018; Accepted 23 September 2018

Available online 24 September 2018

0145-305X/© 2018 Elsevier Ltd. All rights reserved.

opportunistic pathogen of plants, insects, nematodes and human. Several insect species have been used to study *P. aeruginosa* pathogenesis and virulence. These include the greater waxworm *Galleria mellonella* (Jander et al., 2000; Miyata et al., 2003; Andrejko et al., 2009; Koch et al., 2014; Doberenz et al., 2017), fruit fly *Drosophila melanogaster* (Apidianakis et al., 2005; Lutter et al., 2012; Haller et al., 2014; Martínez and Campos-Gómez, 2016), silkworm (Kaito et al., 2002; Iiyama et al., 2007; Okuda et al., 2010; Chieda et al., 2011), and mealworm *Tenebrio molitor* (Park et al., 2014). In the present study, we attempted to identify secreted virulent factors of *P. aeruginosa* by using larval silkworm as the host. Through biochemical and bacterial mutagenesis studies, we obtained results showing that *P. aeruginosa* secretes a metalloprotease (elastase B) as a virulent factor into the hemolymph to manipulate the host PPO activation and AMP production. Possible molecular mechanisms for its effects on the humoral responses are explored and discussed.

## 2. Materials and methods

### 2.1. Identification of major extracellular proteins secreted by *P. aeruginosa* (PAO1)

The secreted proteins of PAO1 were isolated as described previously (Wu et al., 2008; Altindis et al., 2015) with some modifications. Briefly, PAO1 was cultured in Luria-Bertani medium at 37 °C overnight and cell growth was monitored by taking absorbance readings at 600 nm until the  $A_{600nm}$  reached about 2.3. The cells were removed by centrifugation at 8,000g for 20 min at 4 °C, and the supernatant was filtered through a 0.22 µm pore-size filter. In 100 ml of the filtered supernatant, 20 ml of 100% trichloroacetic acid (TCA) was added and incubated on ice overnight before centrifugation at 16,000 g at 4 °C for 60 min. The resulting protein pellet was washed three times with 100% ice-cold acetone. Then the pellet was resuspended in SDS-PAGE sample buffer, incubated at 95 °C for 5 min, and separated by SDS-PAGE. The major bands were excised and digested using In-Gel Tryptic Digestion Kit (Thermo, USA) and analyzed by ion trap LC-MS/MS on Thermo Scientific LTQ XL.

### 2.2. Purification of *P. aeruginosa* elastase B

The PAO1 culture supernatant was prepared as above, filtered through a 0.22 µm pore-size filter, and concentrated 30 fold by ultrafiltration in a centrifugal device (Amicon Ultra-15, NMWL: 10 kDa, Merck Millipore, America). The concentrate was dialyzed against phosphate buffered saline (PBS, 0.27 g/L  $KH_2PO_4$ , 1.42 g/L  $Na_2HPO_4$ , 8 g/L NaCl and 0.2 g/L KCl) overnight. Proteins were then precipitated by ammonium sulfate at 75% of saturation. After centrifugation at 16,000g for 30 min at 4 °C, the protein pellet was dissolved in 50 mM Tris-HCl, pH 8.3 and dialyzed against the same buffer overnight. The dialyzed elastase B was loaded onto an anion-exchange Q-Sepharose Fast-Flow column (2 ml bed volume, GE Healthcare Life Sciences, Sweden) and eluted sequentially with 3 ml of 50 mM Tris-HCl, pH 8.3, containing 100 mM, 250 mM and 1M NaCl. The fractions containing elastase B were concentrated 5 fold in a centrifugal filter (Amicon Ultra-0.5, NMWL: 10 kDa, Merck Millipore, America), dialyzed against 50 mM Tris-HCl, pH 8.3, and fractionated on the Q-Sepharose column under the same condition. The concentrated elastase B fractions were loaded onto an HPLC gel filtration column (TSKgel SuperSW2000, TOSOH, Japan) running with 50 mM sodium phosphate buffer (pH 8.0) containing 150 mM NaCl at a flow rate of 0.5 ml/min. The purified elastase B fractions were pooled, concentrated, dialyzed against PBS, and stored at –80 °C.

**Table 1**  
Oligonucleotide primers used in this study.


Primers	sequence (5'-3')
<b><i>ΔlasB</i></b>	
<i>P. a lasB</i> up F	CCTGGGATCCGGGCTGATCAGCAAGTTC
<i>P. a lasB</i> up R	GGTTGTACACGAGTTTGGACACGTCGATC
<i>P. a lasB</i> down F	GTCCAAACTCGTGTACAACCGTGCCTTC
<i>P. a lasB</i> down R	CCTGAAGCTTGAACCGGTGATGCGATG
<b><i>ΔlasB-C</i></b>	
<i>P. a lasB</i> full F	CCTGAAGCTTGCCTCGGCCGAGTACTTC
<i>P. a lasB</i> full R	CCTGGGATCCGAGCTTACAACGCGCTCG
<b><i>lasB</i> expression</b>	
<i>P. a lasB</i> F	CAACCAGAAGATCGGCAAGTA
<i>P. a lasB</i> R	CGAATTGGCCAACAGGTAGA
<i>P. a GAPDH</i> F	GCAGGTAGTGGCATCAAT
<i>P. a GAPDH</i> R	GGTTCTGGTCGTTGGTGTAG
<b>Quantitative real-time-PCR</b>	
<i>B. m gloverin2</i> F	ACGGACCTTCTGATTACGC
<i>B. m gloverin2</i> R	CATTTCTGTTCCGCCAGT
<i>B. m gloverin3</i> F	GACACGAGAATGGGAGGAG
<i>B. m gloverin3</i> R	AAGACCCTGGTGCCGTAA
<i>B. m gloverin4</i> F	CTTGACAAGAACACCCGCCCT
<i>B. m gloverin4</i> R	GTCTTGAAGGGATCTTCTGGAT
<i>B. m IF4A</i> F	TCTGGCATCATACCTTCTACAA
<i>B. m IF4A</i> R	TCTGTGTCATCTTTCCCTGTT
<i>P. a aprA</i> F	AGTTCCAAGCTGGTGTCTC
<i>P. a aprA</i> R	CCTTCTCGTTGAGGTGTGATCTT
<i>P. a lasA</i> F	CGCTGAATGACGACCTGTT
<i>P. a lasA</i> R	CTTTCGGGTTGATGCTGTAGTA
<i>P. a lepA</i> F	AGGACTGGTCGGATACAGTT
<i>P. a lepA</i> R	TTTACGTTGAGGCCGATGAG
<i>P. a piv</i> F	ATACCCTGACCGTGAAGT
<i>P. a piv</i> R	GGAGTCGGCGAAATACGATAC
<i>P. a GAPDH</i> F(Q)	GCTGGTGTCCGGTGGATTT
<i>P. a GAPDH</i> R(Q)	AGCATGCGGTTGGAGAAG

The restriction enzyme sites *Bam*HI and *Hind*III are underlined.


### 2.3. Preparation of PAO1 *lasB* deletion (*ΔlasB*) and *ΔlasB* complemented (*ΔlasB-C*) mutants

As described previously (Lin et al., 2015), the 866 bp upstream and 668 bp downstream fragments of the *lasB* gene were amplified by overlapping PCR using *Pfu* DNA polymerase and the primer pairs (*lasB* up F/R and *lasB* down F/R) (Table 1). The product was inserted into the *Bam*HI and *Hind*III sites of pK18*mobsacB*, a suicide vector. The gentamicin resistance gene cassette from p34S-Gm was subsequently inserted into the *Hind*III site to yield the mutation plasmid pK-B. After mating an *E. coli* S17-1 derivative carrying pK-B with *P. aeruginosa* (PAO1) on LB plates at 37 °C for 48 h, the cells were suspended in LB at appropriate dilutions and spread on LB plates containing chloramphenicol (to select against the donor strain) and gentamicin (to select for recipient with non-replicating plasmid integrated into its genome). Several colonies were transferred to LB medium and incubated at 37 °C overnight before appropriate dilutions were spread on LB plates containing 12% sucrose for counter-selection against single cross-over mutants. Double cross-over mutants resulting in the nonpolar deletion of *lasB* were verified by PCR using external primer pair *lasB* up F and *lasB* down R and DNA sequencing. The resulting *lasB* deletion mutant contains the first 79 codons fused in frame with the last 65 codons.

To generate *lasB* deletion-complemented mutant, a 1549 bp DNA fragment containing the entire *lasB* opening reading frame was amplified by PCR using *Pfu* DNA-polymerase with *lasB* full F/R primer pairs (Table 1). After DNA sequence verification, the PCR product was inserted into the *Bam*HI and *Hind*III sites of pBBR1MCS-5. The recombinant plasmid was electro-transfected into competent PAO1-*ΔlasB* recipient cells. The complemented strain was selected on gentamicin-kanamycin LB plate and DNA sequence verification.



**Fig. 1. Identification of the PAO1 secreted proteins and purification of elastase B.** (A) The secreted proteins of PAO1 was separated by SDS-PAGE, visualized by Coomassie Brilliant Blue R-250, followed by in-gel trypsin digestion and mass spectrometry analysis. The identified proteins are indicated. Lane M, marker proteins; lane S, PAO1 secreted proteins. (B) Lane 1, fraction collected from 100 mM NaCl elution on the second run of Q-Fast Flow Sepharose column. Lane 2, purified elastase B from the HPLC gel filtration column. The protein bands were visualized by silver staining. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)



**Fig. 2. PAO1 elastase B reduces silkworm phenoloxidase (PO) activity and melanization.** (A) The hemolymph PO activity was measured after injection of purified PAO1 elastase B into the larval silkworm, with phosphate-buffered saline (PBS) and bovine serum albumin (BSA) as controls. (B) Q-Fast Flow Sepharose beads were incubated with purified elastase B or BSA and then injected into the larval hemocoel. The beads were recovered from dissected larvae for melanization inspection under a microscope 4 h later. Each dot represents an individual larva. (C) PO activity of hemolymph from the larvae injected with purified PAO1 elastase B, activated by incubation with curdlan for 30 min. (D) The hemolymph samples were collected from larvae at 24 h post injection of PAO1 elastase B or BSA, activated by curdlan, and then separated on 8% SDS-PAGE followed by immunoblotting using silkworm PPO-1 and -2 antibodies.

#### 2.4. Growth rate assay of PAO1 strains *in vitro*

To compare the growth rate of different strains of PAO1 *in vitro*, the bacteria were cultured till the  $A_{600nm} = 0.6$ . 400  $\mu$ l of these suspensions were added into 50 ml of Luria-Bertani medium respectively and cultured at 37 °C. The  $A_{600}$  was measured by spectrophotometer every 2 h.


#### 2.5. Silkworm rearing and bacteria preparation

The silkworm *B. mori* (Nistari strain) was reared on fresh mulberry leaves at 27 °C in 70% relative humidity and a photoperiod of 13:11 (light: dark). Day 3, 5th instar silkworm larvae were used in all experiments. The PAO1 wide type (WT), *lasB* deletion mutant ( $\Delta$ *lasB*),  $\Delta$ *lasB* complemented ( $\Delta$ *lasB-C*), and  $\Delta$ *lasB-C* control ( $\Delta$ *lasB-pBBR1MCS-5*) strains were respectively cultured at 37 °C in LB medium till  $A_{600nm}$

reached 0.8. The bacteria cells were collected by centrifugation at 8,000g for 20 min at 4 °C. The cells were washed for three times with sterilized PBS and resuspended to  $1 \times 10^8$  CFU/50  $\mu$ l.

#### 2.6. Phenoloxidase (PO) activity assay and prophenoloxidase (PPO) cleavage

After anesthetization on ice for 20 min, day 3, 5th instar silkworm larvae were injected with 10  $\mu$ l of PBS, 10  $\mu$ l of 30  $\mu$ g/ml bovine serum albumin (BSA, BIO BASIC, Canada) in PBS 10  $\mu$ l of 30  $\mu$ g/ml purified elastase B or 50  $\mu$ l of PBS, and 50  $\mu$ l bacterial preparations ( $1 \times 10^8$  CFU/50  $\mu$ l PBS). At different time points, the hemolymph was collected and centrifuged immediately at 16,000g for 30 s at 4 °C to remove hemocytes and bacterial cells. Aliquots (2  $\mu$ l) of plasma sample were added to wells of a 96-wells plate, 100  $\mu$ l of 2 mM dopamine in



**Fig. 3.** PAO1 elastase B induces the production of silkworm gloverins. (A) The silkworm hemolymph samples from larvae at 24 h after injection of PAO1 elastase B, PBS and BSA were subjected to bacterial growth inhibition assays. PAO1 was used as the indicating bacterium on the plates. The data was from five individual larvae. (B) Boiled hemolymph samples from the silkworm larvae 24 h post injection were separated on a 16% Tricine-SDS-PAGE and the bands were digested with trypsin and analyzed by mass spectrometry. The band corresponded to silkworm gloverin is indicated by an arrow. (C) Gloverin-2, -3, and -4 relative mRNA levels in the fat body collected from larvae at 12 h post injection were determined by quantitative real-time PCR. The samples from PBS-injected larvae were used as control. All analyses were performed with five biological replicates.

50 mM sodium phosphate was then added and mixed immediately. The absorbance at 490 nm was recorded every 30 s for 15 min on a microplate reader (Tecan, Pro200, Switzerland), and the maximum slope was determined.

Fifty µl of fresh plasma samples were collected as described above, added simultaneously to 1.5 ml Eppendorf tubes that contained 2 µl of 1 mg/ml curdlan from *Alcaligenes faecalis* (InvivoGen, USA), and incubated at room temperature for 30 min. These mixed samples (2 µl) were taken to measure the PO activity as described above.

For PPO activation assay in hemolymph of protein- or bacteria-injected silkworm, 50 µl aliquots of the plasma samples freshly collected from protein-injected larvae at 24 h or bacteria-injected at 6 h post injection were transferred to 1.5 ml Eppendorf tubes containing 2 µl of 1 mg/ml curdlan and incubated at room temperature for 30 min. Then, each sample was mixed with 52 µl of 2 × SDS loading buffer and boiled immediately at 100 °C for 5 min. After centrifugation at 10,000g for 3 min, the samples were separated by 8% SDS-PAGE, electro-transferred onto PVDF membrane, and subjected to immunoblot analysis using 1:10,000 diluted PPO1 and PPO2 antisera (gifts from Dr. Michael Strand) as the primary antibodies. The proteolytic processing of PPO1 and PPO2 to PO1 and PO2 was visualized using western blotting detection kit (Advansta, USA) on a Chemiluminescence imaging system (ChemiScope Mini2950, Clinx, China).

## 2.7. Melanotic encapsulation of Sepharose beads

Encapsulation of Sepharose beads in silkworm hemolymph was carried out as described before (Li et al., 2016). The QAE Sepharose Fast Flow chromatography beads (50–150 µm in diameter, Sigma, USA) were fully stained with 0.1% Congo-Red for 2 h and then washed three times with 50 mM Tris-HCl (pH 8.3). The washed beads (50 µl

containing 100 beads) were incubated in 1 ml of purified elastase B or BSA (6 µg/ml in 50 mM Tris-HCl, pH 8.3) for 1 h at room temperature with gentle rotation. Fifty µl aliquots of the bead suspensions were injected into larval hemocoel. After 4 h, the larvae were dissected to recover the beads for observation under microscope.


## 2.8. Expression analysis of PAO1 *lasB* gene in the hemocoel of *B. mori*

Bacteria preparations of wild-type,  $\Delta lasB$ ,  $\Delta lasB-C$  and  $\Delta lasB$ -pBBR1MCS-5 PAO1 cells were separately injected into hemocoel of *B. mori* larvae. Total RNA was extracted from the PAO1 bacteria cells and hemocytes harvested from hemolymph of the infected silkworm at 3, 6 and 12 hpi using TriPure Isolation Reagent (Roche, Basel, Switzerland) and purified by Direct-zol™ RNA Miniprep (Zymo Research, California, USA). cDNA was synthesized from the purified RNA using Transcriptor First Strand cDNA Synthesis Kit (Roche, Basel, Switzerland) with random hexamers. Resulting cDNA was used as a template for PCR amplification by *lasB* F/R primer pair with *GAPDH* as an internal control, and the primers used in this experiment were listed in Table 1.

## 2.9. Elastolytic activity assay

Elastolytic activity of elastase B produced by PAO1s in hemolymph of PAO1-infected silkworm was determined in an ECR (Elastin-Congo-Red, Sigma, USA) assay (Pearson et al., 1997) with some modifications. At 4, 8, 12 and 24 h post infection, fresh hemolymph samples were quickly collected into tubes containing 10 µl of 1% 1-phenyl-2-thiourea (PTU) per 100 µl hemolymph. The samples were centrifuged at 10,000 g at 4 °C for 1 min and the supernatants were filtered through 0.22 µm pore-size membrane to remove hemocytes and bacteria. Fifty µl plasma was added to tubes containing 20 mg of ECR and 1 ml of





**Fig. 4.** PAO1 *lasB* expresses in the hemolymph of silkworm larvae after infection. (A) The growth curves of different PAO1 strains in Luria-Bertani broth. WT, PAO1 wild type;  $\Delta lasB$ , *lasB* deletion mutant;  $\Delta lasB-C$ , *lasB* complemented; *lasB*-pBBR1MCS-5,  $\Delta lasB-C$  control. (B) PAO1 *lasB* expression after injection into the larval silkworm. The gene expression was analyzed by semi-quantitative PCR with *GAPDH* acting as an internal control. (C) Elastolytic activity in the hemolymph of PAO1 infected-silkworm larvae. The dash lines represent activity in the hemolymph after injection of purified elastase B.

100 mM Tris-HCl, 1 mM CaCl<sub>2</sub> and incubated at 37 °C for 24 h with rotation. Ten  $\mu$ l of 100 mM EDTA was added to stop the reaction. Insoluble ECR was removed by centrifugation, and A<sub>495nm</sub> was measured on a spectrophotometer, corrected by subtracting the A<sub>495nm</sub> of each sample that had been incubated in the absence of ECR. The assay was repeated independently with three biological replications.

#### 2.10. Antimicrobial activity assay and antibacterial peptide genes expression analysis

Hemolymph from five silkworm larvae was collected at 24 h after injection of PBS, BSA and elastase B, or 12 h and 24 h after injection of PAO1. After heating at 100 °C for 5 min, the supernatant was harvested after centrifugation. Zone of inhibition in a thin layer of LB agar plate assay was used to determine antibacterial activity of these samples as described previously (Hoffmann et al., 1981). In brief, 2  $\mu$ l wild type PAO1 culture suspension in log-phase (A<sub>600nm</sub>  $\approx$  0.8) was mixed with 10 ml melt LB agar and poured into sterile Petri dishes. Holes with a diameter of 1.5 mm were made using a cut 1 ml pipet tip. Two  $\mu$ l of the supernatants were added to each hole and the zone of inhibition diameter was measured after overnight incubation at 37 °C. The supernatants were also treated with 4  $\times$  SDS loading buffer at 100 °C for 5 min and separated on a 16% Tricine-SDS- PAGE (Schagger, 2006). The bands were excised and digested using In-Gel Tryptic Digestion Kit (Thermo, USA), and subjected to ion trap LC-MS/MS analysis on Thermo Scientific LTQ.

For AMP genes expression analysis, fat body tissues from five silkworm larvae were collected at 6, 12 and 24 h after injection of PBS, BSA and elastase B, or 12 h after injection of PAO1. Total RNA of the collected fat body was extracted and expression of AMP genes was analyzed by quantitative real-time PCR performed on a Rotor Q (Qiagen, Hilden, Germany) with *IF4A* as an internal control. The primers were

listed in Table 1. The experiment was performed with three biological replicates and the results were calculated using a relative quantitative method ( $2^{-\Delta\Delta Ct}$ ).

#### 2.11. Determination of colony forming units (CFU) of PAO1 in the hemolymph of silkworm and silkworm survival assay

Day 3, 5th instar silkworm larvae were injected with freshly bacteria preparations. After 12, 24 and 36 h, hemolymph of 8 infected larvae from each group was collected in 1.5 ml Eppendorf tubes containing 2  $\mu$ l of 1% PTU on ice. The samples were serially diluted with sterile PBS and spread on LB agar plate containing kanamycin sulfate (50  $\mu$ g/ml). Colonies on the plates were counted after overnight incubation at 37 °C.

Day 3, 5th instar larvae were anaesthetized on ice for 20 min and injected with 50  $\mu$ l fresh bacteria preparations. Each group consisted of at least twenty larvae and survival was monitored every 12 h.


#### 2.12. Statistical analyses

Graphpad Prism 5.0 was used to plot the data. The student's test was used to analyze the statistical significance of PO activity, beads melanization assay, antimicrobial activity, CFUs and the fold change of gene transcripts. Log-rank (Mantel-Cox) test was used to analyze statistical significance.

### 3. Results

#### 3.1. Elastase B from PAO1 secretion suppresses PPO activation but stimulates AMP production in *B. mori*

To identify the proteins PAO1 secretes, we collected the conditioned



**Fig. 5.** Deletion of *lasB* causes an increase in PO activity of hemolymph from PAO1 infected silkworm larvae. (A) PO activity of the hemolymph from larvae injected with PAO1 cells. (B) PO activity of hemolymph collected from larvae injected with PAO1 cells and activated by curdlan. (C) Melanization of hemolymph samples collected from larvae at 6 and 12 h after injection with PAO1 cells and activated by curdlan. These samples were placed on bench and photographed immediately (0 min) after activation and 30 min later. (D) Proteolytic processing of PPO in the hemolymph at 6 h post infection. The hemolymph samples were collected from larvae 6 h post injection with PAO1 cells and activated by curdlan. The samples were separated by 8% SDS-PAGE and visualized by immunoblotting with silkworm PPO antibodies.

culture medium and separated the TCA-precipitated proteins by SDS-PAGE. The major bands on the gel were identified as catalase, aminopeptidase, chitin-binding protein, elastase B and Las A by MALDI-TOF/TOF (Fig. 1A). Elastase B is the most abundant protein in the secretion. After separation by ion exchange and gel filtration chromatography, we obtained about 120 μg of purified elastase B (Fig. 1B) from 3 L of the PAO1 culture broth.


In order to investigate the effect of elastase B on the silkworm immune system, we injected the purified elastase B into the hemolymph of the tested larvae. The hemolymph samples were collected and subjected to PPO activation and PO activity assays. Within 24 h after injection of elastase B, PO activity decreased in an elastase B dosage-dependent manner (Fig. 2A). Melanized beads in the elastase B-injected larvae were fewer than those in the BSA-injected larvae (Fig. 2B), suggesting that elastase B had suppressed melanotic encapsulation. To examine the cleavage activation of PPO, we incubated the hemolymph samples with curdlan. PO activity of the curdlan-activated hemolymph was lower after elastase B treatment (Fig. 2C). Immunoblot results revealed that cleavage of PPO1 and PPO2 was hindered by elastase B (Fig. 2D). These data together indicate that PAO1 elastase B somehow inhibits the silkworm PPO activation.

Along with PPO activation and melanization, induced production of antimicrobial peptides is another immune response. Therefore, we measured antibacterial activity of the silkworm hemolymph. In the samples from larvae that were injected with elastase B, we observed higher antibacterial activity than in the samples from PBS- and BSA-injected larvae (Fig. 3A). This observation prompted us to examine the change of AMP protein levels in the hemolymph after elastase B injection. We separated low molecular weight protein fraction of the heated hemolymph by Tricine-SDS-PAGE and detected a new band at about 16 kDa in the elastase B-injected hemolymph samples, and a


higher dosage of the enzyme resulted in a higher intensity of the band (Fig. 3B). Mass spectrometry identified this band as silkworm gloverins (data not shown). Quantitative PCR results confirmed that gloverin genes expression in the silkworm fat body were up-regulated after elastase B injection (Fig. 3C). All these data suggest that PAO1 elastase B somehow stimulates silkworm AMP production.

### 3.2. PAO1 secretes elastase B into the silkworm hemolymph after invasion

To validate above the results obtained using the purified elastase B, we made a deletion in the elastase B gene (*ΔLasB*) and its corresponding complemented (*ΔLasB-C*) PAO1 strains. Elastolytic activity assays confirmed our successful construction of the deletion and complemented strains (Supplemental Fig. 1). These strains grew in LB medium with no apparent difference (Fig. 4A), indicating that the deletion of *lasB* does not impair the growth of PAO1. The deletion did not affect expression of PAO1 extracellular proteases *aprA*, *lasA*, *lepA*, and *protease IV* (Supplemental Fig. 2). We detected the expression of *lasB* after injection of wild-type and *lasB* complemented PAO1 cells into the hemolymph of the silkworm. Our results demonstrated that PAO1 *lasB* is expressed shortly (3 h) in the silkworm after injection (Fig. 4B). Since the PAO1 elastase B antibody is not available for detection of elastase B protein, we measured elastolytic activity in the silkworm hemolymph after bacterial injection. A rapid increase of the activity was observed at 4 h after infection by the wild type (WT) and *ΔLasB-C* PAO1 (Fig. 4C). Of note, at 12 h after injection of WT, the elastolytic activity was equivalent to that in hemolymph from silkworm that had been injected with 0.3 μg purified elastase B. This assured consistency of our data obtained from live bacterial cells and purified elastase B protein.



**Fig. 6.** Deletion of *lasB* decreases the expression of the silkworm gloverin genes after PAO1 infection. (A) Anti-PAO1 activity of the hemolymph from PAO1-infected larval silkworm. (B) Comparison of the gloverin levels in the hemolymph from PAO1-infected larvae at 24 hpi after Tricine-SDS-PAGE. The band corresponded to gloverin was confirmed by mass spectrometry. (C) Quantitative real-time PCR analysis of expression of *gloverin-2* and *-3* in fat body from larvae at 12 h post infection.



**Fig. 7.** Deletion of *lasB* reduces the pathogenicity of PAO1 to the silkworm larvae. (A) Survival curve of the silkworm larvae infected with  $1 \times 10^8$  CFU PAO1 per larva. (B) CFU of PAO1 in hemolymph from the infected larvae.

**3.3. Deletion of *lasB* results in partial loss of the ability of PAO1 in PPO activation inhibition and to AMP induction in the silkworm**

We next examined PO activity and PPO activation in the hemolymph collected from PAO1 infected larval silkworm. Compared with WT infected, *ΔlasB* infected samples had higher PO activity with (Fig. 5A) or without activation by curdlan (Fig. 5B), suggesting that PAO1 elastase B contributes to the inhibition of PO activity in the silkworm. After 6 h infection, *ΔlasB*-infected samples showed obvious melanization but WT-infected sample did not (Fig. 5C). Complementation of *ΔlasB* partially restored the inhibition of PO activity and melanization (Fig. 5A, B and 5C), implying that other PAO1 factors contribute to the host PPO activation inhibition along with elastase B during the infection. Immunoblotting revealed the proteolytic activation of PPO in samples from larvae at 6 h post infection with different PAO1 strains (Fig. 5D) was consistent with the PO activity data and hemolymph melanization.

We compared the antimicrobial activity of silkworm hemolymph after PAO1 infection. Overall, PAO1 infection induced strong antibacterial activity in the hemolymph (Fig. 6A). Hemolymph from PAO1 *ΔlasB*-infected larvae displayed slightly lower activity than that from the WT-infected larvae, and PAO1 *ΔlasB-C*-infected hemolymph had an activity between the WT- and *ΔlasB*-infected samples. Gloverin protein levels (Fig. 6B) and mRNA levels (Fig. 6C) in the silkworm infected with different PAO1 strains exhibited similar patterns as the antibacterial activity. These results confirm that elastase B is one of the factors that induce AMP production when PAO1 infects the silkworm.

**3.4. Deletion of *lasB* results in higher susceptibility of PAO1 to the silkworm defense**

We finally compared the survival of silkworm larvae and proliferation of the bacteria in the silkworm hemolymph after PAO1 infection. We injected  $1 \times 10^8$  cells of the four PAO1 strains into each

larvae at the dosage used in the assays of *lasB* expression, elastolytic activity, silkworm PO activity, PPO activation, and antibacterial activity. No larvae survived over 2.5 days in the group of WT infection, whereas about 5% survived at 4 days in the group of  $\Delta$ *lasB*-infected group (Fig. 7A). Bacterial colony counting indicated that PAO1 propagated fast in the silkworm hemolymph 12 h after injection (Fig. 7B). PAO1  $\Delta$ *lasB* grew slower than the WT in the hemolymph, indicating that deletion of *lasB* caused PAO1 to be less virulent and more susceptible to the host immune reactions. If we pre-injected the silkworm with PO inhibitor 1-phenyl-2-thiourea (PTU), the survival of infected larvae and bacterial CFU in the larval hemolymph had no significant difference between the WT- and *lasB*-infected groups (Supplemental Fig. 3). This result reinforced the critical role of PO in killing of bacterium PAO1 and the potential role of PAO1 elastase B in counteracting of silkworm PO pathway.

#### 4. Discussion

Bacteria produce proteases during invasion and infection of their hosts. For instance, the honey bee pathogenic bacterium *Paenibacillus larvae* produces a metalloprotease in the host midgut after it orally infects the bees (Antúnez et al., 2011). The insect pathogen *P. luminescens* produces an M4 metalloprotease PrtS to induce melanization in the hemolymph of *M. sexta* and *Drosophila* during infection (Held et al., 2007). *Pseudomonas entomophila* produces an alkaline zinc metalloprotease AprA to suppress the IMD pathway of *Drosophila* (Liehl et al., 2006), and cellular and humoral immune responses of bean bug *Riptortus pedestris* (Lee et al., 2018). In the present study, we showed that PAO1 produces an M4 metalloprotease elastase B in the hemolymph of silkworm during infection (Fig. 4B and C). Serralysin, an M12 metalloprotease secreted by *S. marcescens*, may degrade a serine protease homolog, SPH-1, which is involved in melanization and nodulation in the silkworm (Tokura et al., 2014) and thus suppress the immune responses (Ishii et al., 2014a). In mammalian infection models, elastase B enhance the virulence of *P. aeruginosa* strains through directly destructing tissue and damaging cell functions, or indirectly interfering with host-defense (Nicas and Iglewski, 1985; Adekoya and Sylte, 2009). Elastase B protects *P. aeruginosa* cells against phagocytosis by degrading the collectin which is a strong inducer of plant, insect and mammalian immune responses (Samakovlis et al., 1992; Casilag et al., 2015; Hajam et al., 2017; Cui et al., 2018) and it disrupts human vascular cells and endothelial barrier (Beaufort et al., 2011, 2013). Biofilm formation is the predominant growth mode for bacteria in various environments and provides a way to escape host immunity systems (Fux et al., 2003). Elastase B can also increase *P. aeruginosa* attachment, microcolony and biofilm formation to escape host immune responses (Yu et al., 2014). In the greater wax moth larvae, injection of thermolysin, an M4 metalloprotease, induced the PO and antibacterial activities in the hemolymph (Griesch et al., 2000; Altincicek et al., 2007). Similarly, challenge with live *P. aeruginosa* cells or injection with elastase B protein up-regulated PO and antibacterial activities in the larvae of *G. mellonella* (Andrejko and Mizerska-Dudka, 2011; 2014). A recent study in *Drosophila* showed that a *P. aeruginosa* mutant strain of *lasB* caused less expression of drosomycin, an AMP induced by the Toll pathway (Issa et al., 2018). In our study, by using the purified protein and mutant bacterial strains, we clearly demonstrated that PAO1 elastase B plays a role in AMP induction but PPO activation suppression in the silkworm. Regarding the effect of metalloproteases on the insect PPO activation system, we speculate that the difference between the results from us and others might arise from the differences in bacteria strains (Andrejko et al., 2014), hosts, and experimental procedures.

AMP generation and PO-catalyzed melanization are two critical defense responses of insects. In general, AMP gene transcription is mediated through the Toll or IMD pathways upon recognition of invading pathogens or parasites by the host. Activation of the Toll pathway and PPO is mediated by an extracellular clip-domain serine

protease cascade (Kambris et al., 2006; Tang et al., 2006; An et al., 2009; Kanost and Jiang, 2015), which is tightly regulated by serine protease inhibitors in the serpin superfamily (Meekins et al., 2016). To find how the PAO1 elastase B affects silkworm PPO activation and AMP production, we examined proteins in the elastase B-injected hemolymph of larval silkworm and tobacco hornworm (Supplemental Fig. 4). Injection of elastase B resulted in decrease of silkworm inter-alpha-trypsin inhibitor H4-like protein, hemocyte aggregation inhibitor protein (HAIP), serpin-1A and serpin-9 (Supplemental Fig. 4A). Inter-alpha-trypsin inhibitor plays a major role in extracellular matrix stability and integrity (Bost et al., 1998). In our study, we found that a high dosage of elastase B caused a breakup of the silkworm gut tissue (data not shown). Human vascular cells and endothelial barrier were also susceptible to elastase B (Beaufort et al., 2011, 2013). HAIPs were suggested to serve in *Drosophila* hemolymph clotting and epithelial defense against pathogens (Kucerova et al., 2015; Pesch et al., 2015). Among the decreased proteins, serpin-1A and -9 are of particular interest. We previously identified serpin-5 as a negative regulator of PPO activation and AMP induction pathways in the silkworm (Li et al., 2016). We demonstrated in this study that injection of elastase B resulted in a decrease of serpin-1A and -9 and an increase of AMP production. This implies that these two serpins might be involved in regulation of silkworm AMP induction. To take the advantage of the antibodies against serine proteases, serine protease homologs (SPHs) and serpins in *M. sexta* hemolymph, we injected elastase B into hemocoel of the tobacco hornworm and examined the changes of hemolymph protein by immunoblotting (Supplemental Fig. 4B). Injection of elastase B into tobacco hornworm strongly inhibited PO activity in the hemolymph (Supplemental Figs. 5A and 5B), as that did to the silkworm (Fig. 2). Among about twenty proteins we examined, activation of SPH-1, -2 and -33 were somewhat hampered by elastase B (Supplemental Fig. 4B). The interactions among SPH-1/2, PPOs and PPO-activating protease (PAP) known to be required for generating active PO (Yu et al., 2003; Gupta et al., 2005). It is possible that elastase B impaired SPH-1 and -2 cleavage activation by an unknown mechanism and thus reduced hemolymph PPO activation in the hornworm. Given the finding that silkworm SPH-1 is involved in melanization (Tokura et al., 2014), it is possible that elastase B suppresses silkworm PPO activation in a similar manner.

It has been reported that another protease secreted by PAO1, protease IV, interferes with AMP production but activates PPO and that elastase B induces generation of AMPs in the mealworm (Park et al., 2014). We found that injection of elastase B into *Tenebrio molitor* larvae strongly reduced PPO activation (Supplemental Fig. 5C), as observed in the silkworm and tobacco hornworm. These results raise an interesting question: why a simple bacterium needs different secreted proteases to modulate the immune pathways of its insect host in opposite ways? As a matter of fact, bacteria produce an array of proteases as virulent factors to interfere with host defense responses. These proteases may destruct components of the immune pathways, interfere intracellular immune signaling, inactivate AMPs, impair phagocytosis, and so on (Potempa and Pike, 2009). From *P. aeruginosa* KU2 strain, a large exoprotease (LepA) was shown to activate NF- $\kappa$ B-driven promoters (Kida et al., 2008). LepA acts in concert with hemolytic phospholipase C (PlcH) for the virulence and growth of KU2 in mice (Kida et al., 2011). A recent work showed that, in addition to alkaline protease AprA, elastase B can degrade exogenous flagellin to prevent flagellin-mediated immune recognition. It is thought that production of the two proteases with flagellin-degrading activity provides an infallible mechanism for *P. aeruginosa* to evade recognition by the host (Casilag et al., 2015). The findings in our current study and made by others (Park et al., 2014) suggest that a more complicated scenario about the interplay of bacterial virulent factors during invasion and infection of the host.

In summary, we presented evidence that PAO1 secretes elastase B as a virulent factor during its infection of a lepidopteran insect, *B. mori*. Elastase B inhibits PPO activation but somehow induces AMP



production in the silkworm hemolymph. Our study suggests more complex interactions between virulent factors of bacterial pathogens and host immune systems.

#### Declaration of interest statement

The authors declare that there are no conflict of interest.

#### Acknowledgment

This study was supported by the National Science Foundation of China under grants 31272497 and 31772530 to ZL; and by the National Institutes of Health under grants AI12662 and GM58634 to HJ. The authors would like to thank Dr. Michael Strand at University of Georgia for providing the antisera to the silkworm PPO-1 and -2.

#### Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.dci.2018.09.017>.

#### References

- Adekoya, O.A., Sylte, I., 2009. The thermolysin family (M4) of enzymes: therapeutic and biotechnological potential. *Chem. Biol. Drug Des.* 73, 7–16.
- Altincek, B., Linder, M., Linder, D., Preissner, K.T., Vilcinskis, A., 2007. Microbial metalloproteinases mediate sensing of invading pathogens and activate innate immune responses in the lepidopteran model host *Galleria mellonella*. *Infect. Immun.* 75, 175–183.
- Altindis, E., Dong, T., Catalano, C., Mekalanos, J., 2015. Secretome analysis of *Vibrio cholerae* type VI secretion system reveals a new effector-immunity pair. *mBio* 2, e00075.
- An, C., Ishibashi, J., Ragan, E.J., Jiang, H., Kanost, M.R., 2009. Functions of *Manduca sexta* hemolymph proteinases HP6 and HP8 in two innate immune pathways. *J. Biol. Chem.* 284, 19716–19726.
- Andrejko, M., Mizerska-Dudka, M., 2011. Elastase B of *Pseudomonas aeruginosa* stimulates the humoral immune response in the greater wax moth, *Galleria mellonella*. *J. Invertebr. Pathol.* 107, 16–26.
- Andrejko, M., Mizerska-Dudka, M., Jakubowicz, T., 2009. Antibacterial activity in vivo and in vitro in the hemolymph of *Galleria mellonella* infected with *Pseudomonas aeruginosa*. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* 152, 118–123.
- Andrejko, M., Zdybicka-Barabas, A., Cytirynska, M., 2014. Diverse effects of *Galleria mellonella* infection with entomopathogenic and clinical strains of *Pseudomonas aeruginosa*. *J. Invertebr. Pathol.* 115, 14–25.
- Antúnez, K., Arredondo, D., Anido, M., Zunino, P., 2011. Metalloprotease production by *Paenibacillus larvae* during the infection of honeybee larvae. *Microbiology* 157, 1474–1480.
- Apidianakis, Y., Mindrinos, M.N., Xiao, W., Lau, G.W., Baldini, R.L., Davis, R.W., Rahme, L.G., 2005. Profiling early infection responses: *Pseudomonas aeruginosa* eludes host defenses by suppressing antimicrobial peptide gene expression. *Proc. Natl. Acad. Sci. U.S.A.* 102, 2573–2578.
- Beaufort, N., Corvazier, E., Hervieu, A., Choqueux, C., Dussiot, M., Louedec, L., Cady, A., de Bentzmann, S., Michel, J.B., Pidard, D., 2011. The thermolysin-like metalloproteinase and virulence factor LasB from pathogenic *Pseudomonas aeruginosa* induces anoikis of human vascular cells. *Cell Microbiol.* 13, 1149–1167.
- Beaufort, N., Corvazier, E., Mlanaoindrou, S., Bentzmann, S.D., Pidard, D., 2013. Disruption of the endothelial barrier by proteases from the bacterial pathogen *Pseudomonas aeruginosa*: implication of matrilysin and receptor cleavage. *PLoS One* 8, e75708.
- Bost, F., Diarra-Mehrpour, M., Martin, J.P., 1998. Inter- $\alpha$ -trypsin inhibitor proteoglycan family. *Eur. J. Biochem.* 252, 339–346.
- Casilag, F., Lorenz, A., Krueger, J., Klawonn, F., Weiss, S., Häussler, S., 2015. The LasB elastase of *Pseudomonas aeruginosa* acts in concert with alkaline protease AprA to prevent Flagellin-mediated immune recognition. *Infect. Immun.* 84, 162–171.
- Chieda, Y., Iiyama, K., Lee, J.M., Kusakabe, T., Yasunaga-Aoki, C., Shimizu, S., 2011. Virulence of an exotoxin A-deficient strain of *Pseudomonas aeruginosa* toward the silkworm, *Bombyx mori*. *Microb. Pathog.* 51, 407–414.
- Cui, B., Liu, X., Fang, Y., Zhou, P., Zhang, Y., Wang, Y., 2018. Flagellin as a vaccine adjuvant. *Expert Rev. Vaccines* 17, 335–349.
- Doberenz, S., Eckweiler, D., Reichert, O., Jensen, V., Bunk, B., Spröer, C., Kordes, A., Frangipani, E., Luong, K., Korlach, J., Heeb, S., Overmann, J., Kaever, V., Haussler, S., 2017. Identification of a *Pseudomonas aeruginosa* PAO1 DNA methyltransferase, its targets, and physiological roles. *mBio* 8, e02312–e02316.
- Eleftherianos, I., Boundy, S., Joyce, S.A., Aslam, S., Marshall, J.W., Cox, R.J., Simpson, T.J., Clarke, D.J., French-Constant, R.H., Reynolds, S.E., 2007. An antibiotic produced by an insect-pathogenic bacterium suppresses host defenses through phenoloxidase inhibition. *Proc. Natl. Acad. Sci. U.S.A.* 104, 2419–2424.
- Fux, C.A., Stoodley, P., Hall-Stoodley, L., Costerton, J.W., 2003. Bacterial biofilms: a diagnostic and therapeutic challenge. *Expert Rev. Anti Infect. Ther.* 1, 667–683.
- Griesch, J., Wedde, M., Vilcinskis, A., 2000. Recognition and regulation of metalloproteinase activity in the haemolymph of *Galleria mellonella*: a new pathway mediating induction of humoral immune responses. *Insect Biochem. Mol. Biol.* 30, 461–472.
- Gupta, S., Wang, Y., Jiang, H., 2005. *Manduca sexta* prophenoloxidase (proPO) activation requires proPO-activating proteinase (PAP) and serine proteinase homologs (SPHs) simultaneously. *Insect Biochem. Mol. Biol.* 35, 241–248.
- Hajam, I.A., Dar, P.A., Shah Nawaz, I., Jaume, J.C., Lee, J.H., 2017. Bacterial flagellin—a potent immunomodulatory agent. *Exp. Mol. Med.* 49, e373.
- Haller, S., Limmer, S., Ferrandon, D., 2014. Assessing *Pseudomonas* virulence with a nonmammalian host: *Drosophila melanogaster*. *Meth. Mol. Biol.* 1149, 723–740.
- Held, K.G., LaRock, C.N., D'Argenio, D.A., Berg, C.A., Collins, C.M., 2007. A metalloprotease secreted by the insect pathogen *Photographus luminescens* induces melanization. *Appl. Environ. Microbiol.* 73, 7622–7628.
- Hoffmann, D., Hultmark, D., Boman, H.G., 1981. Insect immunity: *Galleria mellonella* and other lepidoptera have cecropin-P9-like factors active against gram-negative bacteria. *Insect Biochem.* 11, 537–548.
- Huang, L., Cheng, T., Xu, P., Cheng, D., Fang, T., Xia, Q., 2009. A genome-wide survey for host response of silkworm, *Bombyx mori* during pathogen *Bacillus bombysepticus* infection. *PLoS One* 4, e8098.
- Hurst, G.D., Anbutu, H., Kutsukake, M., Fukatsu, T., 2003. Hidden from the host: *Spiroplasma* bacteria infecting *Drosophila* do not cause an immune response, but are suppressed by ectopic immune activation. *Insect Mol. Biol.* 12, 93–97.
- Iiyama, K., Chieda, Y., Lee, J.M., Kusakabe, T., Yasunaga-Aoki, C., Shimizu, S., 2007. Effect of superoxide dismutase gene inactivation on virulence of *Pseudomonas aeruginosa* PAO1 toward the silkworm, *Bombyx mori*. *Appl. Environ. Microbiol.* 73, 1569–1575.
- Ishii, K., Adachi, T., Hamamoto, H., Sekimizu, K., 2014a. *Serratia marcescens* suppresses host cellular immunity via the production of an adhesion-inhibitory factor against immunosurveillance cells. *J. Biol. Chem.* 289, 5876–5888.
- Ishii, K., Adachi, T., Hara, T., Hamamoto, H., Sekimizu, K., 2014b. Identification of a *Serratia marcescens* virulence factor that promotes hemolymph bleeding in the silkworm, *Bombyx mori*. *J. Invertebr. Pathol.* 117, 61–67.
- Ishii, K., Adachi, T., Imamura, K., Takano, S., Usui, K., Suzuki, K., Hamamoto, H., Watanabe, T., Sekimizu, K., 2012. *Serratia marcescens* induces apoptotic cell death in host immune cells via a lipopolysaccharide- and flagella-dependent mechanism. *J. Biol. Chem.* 287, 36582–36592.
- Issa, N., Guillaumont, N., Lauret, E., Matt, N., Schaeffer-Reiss, C., Van Dorselaer, A., Reichhart, J.M., Veillard, F., 2018. The circulating protease periphonse is an immune sensor for microbial proteolytic activities upstream of the *Drosophila* Toll pathway. *Mol. Cell* 69, 539–550 e6.
- Jander, G., Rahme, L.G., Ausubel, F.M., 2000. Positive correlation between virulence of *Pseudomonas aeruginosa* mutants in mice and insects. *J. Bacteriol.* 182, 3843–3845.
- Kaito, C., Akimitsu, N., Watanabe, H., Sekimizu, K., 2002. Silkworm larvae as an animal model of bacterial infection pathogenic to humans. *Microb. Pathog.* 32, 183–190.
- Kambris, Z., Brun, S., Jang, I.H., Nam, H.J., Romeo, Y., Takahashi, K., Lee, W.J., Ueda, R., Lemaitre, B., 2006. *Drosophila* immunity: a large-scale in vivo RNAi screen identifies five serine proteases required for Toll activation. *Curr. Biol.* 16, 808–813.
- Kanost, M.R., Jiang, H., 2015. Clip-domain serine proteases as immune factors in insect hemolymph. *Curr. Opin. Insect Sci.* 11, 47–55.
- Kida, Y., Higashimoto, Y., Inoue, H., Shimizu, T., Kuwano, K., 2008. A novel secreted protease from *Pseudomonas aeruginosa* activates NF- $\kappa$ B through protease-activated receptors. *Cell Microbiol.* 10, 1491–1504.
- Kida, Y., Shimizu, T., Kuwano, K., 2011. Cooperation between LepA and PlcH contributes to the in vivo virulence and growth of *Pseudomonas aeruginosa* in mice. *Infect. Immun.* 79, 211–219.
- Koch, G., Nadal-Jimenez, P., Cool, R.H., Quax, W.J., 2014. Assessing *Pseudomonas* virulence with nonmammalian host: *Galleria mellonella*. *Meth. Mol. Biol.* 1149, 681–688.
- Kucerova, L., Broz, V., Arefin, B., Maaroufi, H.O., Hurychova, J., Strnad, H., Zurovec, M., Theopold, U., 2015. The *Drosophila* Chitinase-like protein IDGF3 is involved in protection against nematodes and in wound healing. *J. Innate Immun.* 8, 199–210.
- Lee, S.A., Jang, S.H., Kim, B.H., Shibata, T., Yoo, J., Jung, Y., Kawabata, S.I., Lee, B.L., 2018. Insecticidal activity of the metalloprotease AprA occurs through suppression of host cellular and humoral immunity. *Dev. Comp. Immunol.* 81, 116–126.
- Li, J., Ma, L., Lin, Z., Zou, Z., Lu, Z., 2016. Serpin-5 regulates phenoloxidase activation and antimicrobial peptide pathways in the silkworm, *Bombyx mori*. *Insect Biochem. Mol. Biol.* 73, 27–37.
- Liehl, P., Blight, M., Vodovar, N., Bocard, F., Lemaitre, B., 2006. Prevalence of local immune response against oral infection in a *Drosophila/Pseudomonas* infection model. *PLoS Pathog.* 2, e56.
- Lin, J., Cheng, J., Chen, K., Guo, C., Zhang, W., Yang, X., Ding, W., Ma, L., Wang, Y., Shen, X., 2015. The icmF3 locus is involved in multiple adaptation- and virulence-related characteristics in *Pseudomonas aeruginosa* PAO1. *Front. Cell. Infect. Microbiol.* 5, 70.
- Lutter, E.I., Purighalla, S., Duong, J., Storey, D.G., 2012. Lethality and cooperation of *Pseudomonas aeruginosa* quorum-sensing mutants in *Drosophila melanogaster* infection models. *Microbiology* 158, 2125–2132.
- Martinez, E., Campos-Gómez, J., 2016. Oxylipins produced by *Pseudomonas aeruginosa* promote biofilm formation and virulence. *Nat. Commun.* 7, 13823.
- Meekins, D.A., Kanost, M.R., Michel, K., 2016. Serpins in arthropod biology. *Semi. Cell Dev. Biol.* 62, 105–119.
- Miyata, S., Casey, M., Frank, D.W., Ausubel, F.M., Drenkard, E., 2003. Use of the *Galleria mellonella* caterpillar as a model host to study the role of the type III secretion system in *Pseudomonas aeruginosa* pathogenesis. *Infect. Immun.* 71, 2404–2413.
- Nicas, T., Iglewski, B.H., 1985. The contribution of exoproducts to virulence of *Pseudomonas aeruginosa*. *Can. J. Microbiol.* 31, 387–392.
- Okuda, J., Hayashi, N., Okamoto, M., Sawada, S., Minagawa, S., Yano, Y., Gotoh, N., 2010. Translocation of *Pseudomonas aeruginosa* from the intestinal tract is mediated

- by the binding of ExoS to an Na,K-ATPase regulator, FXD3. *Infect. Immun.* 78, 4511–4522.
- Park, S.J., Kim, S.K., So, Y.I., Park, H.Y., Li, X.H., Yeom, D.H., Lee, M.N., Lee, B.L., Lee, J.H., 2014. Protease IV, a quorum sensing-dependent protease of *Pseudomonas aeruginosa* modulates insect innate immunity. *Mol. Microbiol.* 94, 1298–1314.
- Pearson, J.P., Pesci, E.C., Iglewski, B.H., 1997. Roles of *Pseudomonas aeruginosa* las and rhl quorum-sensing systems in control of elastase and rhamnolipid biosynthesis genes. *J. Bacteriol.* 179, 5756–5767.
- Pesch, Y.Y., Riedel, D., Patil, K.R., Loch, G., Behr, M., 2015. Chitinases and imaginal disc growth factors organize the extracellular matrix formation at barrier tissues in insects. *Sci. Rep.* 6, 18340.
- Potempa, J., Pike, R.N., 2009. Corruption of innate immunity by bacterial proteases. *J. Innate Immun.* 1, 70–87.
- Sadekuzzaman, M., Park, Y., Lee, S., Kim, K., Jung, J.K., Kim, Y., 2017. An entomopathogenic bacterium, *Xenorhabdus hominickii* ANU101, produces oxindole and suppresses host insect immune response by inhibiting eicosanoid biosynthesis. *J. Invertebr. Pathol.* 145, 13–22.
- Samakovlis, C., Asling, B., Boman, H.G., Gateff, E., Hultmark, D., 1992. In vitro induction of cecropin genes—an immune response in a *Drosophila* blood cell line. *Biochem. Biophys. Res. Commun.* 188, 1169–1175.
- Schagger, H., 2006. Tricine-sds-page. *Nat. Protoc.* 1, 16–22.
- Strand, M.R., 2008. The insect cellular immune response. *Insect Sci.* 15, 1–14.
- Tanaka, H., Ishibashi, J., Fujita, K., Nakajima, Y., Sagisaka, A., Tomimoto, K., Suzuki, N., Yoshiyama, M., Kaneko, Y., Iwasaki, T., Sunagawa, T., Yamaji, K., Asapka, A., Mita, K., Yamakawa, M., 2008. A genome-wide analysis of genes and gene families involved in innate immunity of *Bombyx mori*. *Insect Biochem. Mol. Biol.* 38, 1087–1110.
- Tang, H., Kambris, Z., Lemaitre, B., Hashimoto, C., 2006. Two proteases defining a melanization cascade in the immune system of *Drosophila*. *J. Biol. Chem.* 281, 28097–28104.
- Tokura, A., Fu, G.S., Sakamoto, M., Endo, H., Tanaka, S., Kikuta, S., Tabunoki, H., Sato, R., 2014. Factors functioning in nodule melanization of insects and their mechanisms of accumulation in nodules. *J. Insect Physiol.* 60, 40–49.
- Vallet-Gely, I., Lemaitre, B., Boccard, F., 2008. Bacterial strategies to overcome insect defences. *Nat. Rev. Microbiol.* 6, 302–313.
- Wu, H.Y., Chung, P.C., Shih, H.W., Wen, S.R., Lai, E.M., 2008. Secretome analysis uncovers an Hcp-family protein secreted via a type VI secretion system in *Agrobacterium tumefaciens*. *J. Bacteriol.* 190, 2841–2850.
- Yang, W., Cheng, T., Ye, M., Deng, X., Yi, H., Huang, Y., Tan, X., Han, D., Wang, B., Xiang, Z., Cao, Y., Xia, Q., 2011. Functional divergence among silkworm antimicrobial peptide paralogs by the activities of recombinant proteins and the induced expression profiles. *PLoS One* 6, e18109.
- Yu, H., He, X., Xie, W., Xiong, J., Sheng, H., Guo, S., Huang, C., Zhang, D., Zhang, K., 2014. Elastase LasB of *Pseudomonas aeruginosa* promotes biofilm formation partly through rhamnolipid-mediated regulation. *Can. J. Microbiol.* 60, 227–235.
- Yu, X.Q., Jiang, H., Wang, Y., Kanost, M.R., 2003. Nonproteolytic serine proteinase homologs are involved in prophenoloxidase activation in the tobacco hornworm, *Manduca sexta*. *Insect Biochem. Mol. Biol.* 33, 197–208.